



GFP Expressing Human Dermal Fibroblasts - Neonatal

Catalog #: GFP07

Cell #: > 5X10⁵ cells

Storage: Liquid Nitrogen until ready for culture.
While culturing keep in 37°C CO₂ incubator

Product Format: Cryopreserved

GENERAL INFORMATION

HNDFCs-Neo cells are isolated from normal neonatal foreskin tissue samples and transfected with GFP-Lentiviral particles at passage one. Puromycin resistant GFP-HNDFCs-Neo are selected and shipped in frozen vial (the cell is provided @passage 1). Fibroblast Growth medium (F001, full medium) is recommended for cell culture and these cells have an average minimum population doubling levels > 14 when cultured following the detailed protocol described below). HNDFCs-Neo are tested negative for HIV-1, HBV, HCV, and mycoplasma.

Product is for Research use only

Cells are shipped on dry ice

HANDLING OF ARRIVING CELLS

When you receive the dry ice package with cells in frozen vials, transfer the frozen vials of cells into a -80°C freezer for short period storage or a liquid nitrogen tank for long-term storage.

PROTOCOL FOR THAWING THE CELLS AND SUBCULTURE

Note: If you have any questions or need clarification regarding the protocol for culturing these cells, please reach out to Dr. Jensen Auguste at (978) 608-1766 with your questions before beginning.

1. Pre-coating of T25 flasks- Add 2ml of AlphaBio Coat solution into a T25 flask to cover the whole surface of the flask, 5 mins later, dispose the excessive coating solution by aspiration and the flask is ready to be used.
2. Thaw the frozen cell vial in a 37°C water bath first, and then transfer the cells into the pre-coated T25 flask with 10ml of cAP-44 medium, cells usually become confluent with 1-2 days and ready to be passaged.
3. To passage the cells, rinse the cells in a T25 flask with 5ml HBSS (RT) twice; then add 2ml Cell Detachment solution (RT) into one T25 flask; gently dispose the excessive detachment solution within 20 seconds by aspiration.
4. Leave the T25 flask with the cells at RT or 37°C for 1 min (most cells usually will detach from the surface within 1-2 mins; or monitor the cells under a microscope until most of cells become rounded up, and then gently tap the flask against the bench surface, and the cells will move on the surface of the flask when monitoring under microscope.
5. Add 5ml Trypsin Neutralization Buffer and spin down the cells with 800g centrifugation for 5 mins.

FOR RESEARCH USE ONLY

NEUROMICS REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. v1-09809

www.neuromics.com

Neuromics Antibodies • 5325 West 74th Street, Suite 8 • Edina, MN 55439
Phone 866-350-1500 • fax 612-677-3976 • email: pschuster@neuromics.com

6. Re-suspend the cell pellet with 10 or 15 ml F001 medium and transfer 5 ml each into 2 or 3 pre-coated T25 flasks (for 1/2 to 1/3 subculture ratio).
7. Change medium every 2 or 3 days and the cells usually become confluent within 7 days (when split at a 1/3 ratio).
8. To prepare quiescent cells, when cells are nearly confluent, replace with Fibroblast Basal Medium (F002) containing 0.5%FBS for about 8-12hrs before your experiments.

Note: Should any issues arise while using our cells, our team is here to help troubleshoot any issues. Our cells are backed by our one-time replacement or refund policy. Our recommended protocol including recommended products must be used to be eligible for replacement or refund. Cells that have been refrozen are no longer eligible for refund or replacement.

FOR RESEARCH USE ONLY

NEUROMICS REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. v1-09809

www.neuromics.com

Neuromics Antibodies • 5325 West 74th Street, Suite 8 • Edina, MN 55439
Phone 866-350-1500 • fax 612-677-3976 • email: pshuster@neuromics.com