



## Universal Coating Solution

---

**Catalog #:** AC002

**Product Size:** 20 ml


**Storage:** Frozen or ice packs

**Product Format:** Clear - liquid or frozen

**pH:** 2.5-3.5

**Source:** Human Cells

---



### GENERAL INFORMATION

The biggest challenges for researchers using other coating solutions are a proper coating of the surface, long incubation time of the coating solution, and properly adjusting pH of their vessel once coated. Our product used less time and reduced the risk of killing your cell because of poor coating and unstable PH. We use 6 simple steps:

Universal Coating Solution is easy to use:

1. Thaw solution if frozen
2. Coat your Vessel
3. Incubate for 30 minutes in 37°C incubator
4. Remove excess coating solution
5. Add you cell to the coated vessel
6. Place coated vessel with cell in a 37°C incubator with proper CO<sub>2</sub>, watch it grow

*Product is for Research use only.*

### STORAGE AND USE

All procedures should be done under sterile conditions using aseptic techniques. Avoid prolonged exposure of the protein to ambient temperatures. Repeated freeze/thaw should be avoided. Universal Coating solution is stable for 3 years when stored at -20°C to -80°C. It is stable for at least 3 months when stored at +2°C to +8°C under aseptic conditions. For your convenience, coated plates can be kept for up to 4 weeks when stored aseptically at +2°C to +8°C.

### COATING PROTOCOL

1. If frozen, slowly thaw solution at +2°C to +8°C before use. Undiluted is stable for at least 3 months when stored at +2°C to +8°C under aseptic conditions. Repeated freeze-thaw cycles should be avoided. For longer storage needs, we recommend dividing the thawed stock solution in smaller working aliquots and to store frozen. Frozen stock can be stored up to three years in -20°C to -80°C.

### FOR RESEARCH USE ONLY

NEUROMICS REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. v1-09809

[www.neuromics.com](http://www.neuromics.com)

Neuromics Antibodies • 5325 West 74<sup>th</sup> Street, Suite 8 • Edina, MN 55439  
phone 866-350-1500 • fax 612-677-3976 • e-mail [pshuster@neuromics.com](mailto:pshuster@neuromics.com)


- Guidelines for surface coating calculations can be found in the table below. Lowering the coating concentration might affect the proliferation rate, extending the culture time with about 1 or 2 days. Make sure the coating concentration is high enough to support an even cell growth. For your convenience, coated plates can be kept for up to 4 weeks when stored aseptically at +2°C to +8°C. When culturing cells on the Universal matrices, some cell lines might need an adaptation period and a higher coating concentration is then recommended for the first few passages. Once the cells are adapted to the matrix, the coating concentration usually can be reduced. The coating should be optimized empirically for each cell line and cell type.
- Gently invert the vial to mix the solution. Do not vortex as this may cause fragmentation.

#### GUIDELINE FOR SURFACE COATING

CULTUREWARE	TOTAL COATING SOLUTION VOLUME
6-well	1000 ul/well
12-well	500 ul/well
24-well	300 ul/well
48-well	150 ul/well
96-well	70 ul/well
T25 flask	3000 ul
T75 flask	8000 ul

Do not allow the coated surface to dehydrate as that will inactivate the coating solution. For your convenience, the coated plates can be kept for up to 4 weeks when stored aseptically at +2°C to +8°C. Extra 1xDPBS (Ca<sup>++</sup>/Mg<sup>++</sup>) might have to be added after 1-2 weeks to prevent the plate from drying out.

A panel of different bioassays affirm the media sustain a proper environment for expected cell-type-specific culture, growth, plating, karyotype, physiology, morphology, viability, population doublings, surface markers, cryopreservation, differentiation and/or induction.



#### FOR RESEARCH USE ONLY

NEUROMICS REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. v1-09809

[www.neuromics.com](http://www.neuromics.com)

Neuromics Antibodies • 5325 West 74<sup>th</sup> Street, Suite 8 • Edina, MN 55439  
 phone 866-350-1500 • fax 612-677-3976 • e-mail [pshuster@neuromics.com](mailto:pshuster@neuromics.com)