




<b>Catalog Number:</b>	MO22207	<b>Host:</b>	Mouse
<b>Product Type:</b>	Mouse Monoclonal IgG1	<b>Species Reactivity:</b>	Mouse
<b>Immunogen Sequence:</b>	Recombinant full length annexin A6 expressed in and purified from <i>E. coli</i>	<b>Format:</b>	Purified antibody at 1mg/mL in 50% PBS, 50% glycerol plus 5mM NaN3
<b>Applications:</b>	Immunofluorescence: 1:1,000 Western Blot: 1:1,000-2,000		
<b>Storage:</b>	Dilutions listed as a recommendation. Optimal dilution should be determined by investigator. Antibody can also be aliquotted and stored frozen at -20° C in a manual defrost freezer for six months without detectable loss of activity. The antibody can be stored at 2° - 8° C for 1 month without detectable loss of activity. Avoid repeated freeze-thaw cycles.		

### Application Notes

#### Description/Data

The annexins are a large family of related proteins which share the property of binding to phospholipid containing membranes in a Calcium dependent manner. Annexin A6 has a particularly surprising number of alternate names, 19 being listed on the Genecards ANXA6 site Annexin A6 is normally localized in the cytosol but becomes membrane associated



following cellular injury, and so is often seen in "blebs", transient herniations of the plasma membrane associated with apoptosis and more generally indicative of cell stress. Recent studies suggest that annexin A6 functions, in combination with annexin A4, in the repair of cell membranes following cellular injury.

This antibody was made against full length recombinant human annexin A6 expressed in and purified from *E. coli*. The antibody recognizes annexin A6 in human, rodents and many other mammals and can be used as a marker of blebs, indicative of cellular stress.

*Image: Immunofluorescent analysis of HeLa cells stained with mouse mAb to annexin A6, MO22207, dilution 1:500 in red, and costained with chicken pAb to vimentin, dilution 1:2,000 in green. The blue is Hoechst staining of nuclear DNA. The annexin A6 antibody detects protein predominantly expressed in spherical cellular membrane protrusions, known as blebs, while the vimentin antibody produces strong staining of the intermediate filament network of the cytoskeleton.*

**FOR RESEARCH USE ONLY**

NEUROMICS' REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RISKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE.-V1-/10/2013

[www.neuromics.com](http://www.neuromics.com)

Neuromics Antibodies • 5325 West 74<sup>th</sup> Street, Suite 8 • Edina, MN 55439  
phone 866-350-1500 • fax 612-677-3976 • e-mail: [pshuster@neuromics.com](mailto:pshuster@neuromics.com)