



ELISA Sample Diluents Standard Protocol

Catalog #: KF17352-55

Sizes: 100 ml, 500 ml, 1 L, 10 L

Storage: All of ICT's assay diluents include an antimicrobial agent to provide a useful shelf-life of 1 year when stored at 2-8C.

Sample diluent buffer are used to dilute serum or antigen samples so they read within the functional range of the assay. ICT has created 3 proprietary sample diluent formulations for use with antigen-down and sandwich ELISAs.

Sample Diluent Calculations: simply multiply the value determined from the curve by your dilution factor. For example, if you diluted your sample 1:2 (250uL sample into 250uL sample diluent) and this yielded an OD of 0.437, when read off the curve, the OD of 0.437 correlated to 200 pg/mL, you would simply multiply 200pg/mL by 2 (your dilution factor) yielding a value of 400pg/mL, meaning that the original starting sample had 400pg/mL of analyte present.

FOR RESEARCH USE ONLY

V2-02/13

Neuromics' reagents are for in vitro and certain non-human in vivo experimental use only and not intended for use in any human clinical investigation, diagnosis, prognosis, or treatment. We disclaim all liability in connection with the use of the information contained herein or otherwise, and all such risks are assumed by the user.

phone 866-350-1500 • fax 612-677-3976 • www.neuromics.com