NEUROMICS /



TGF Beta Receptor (type1)

Data Sheet

Catalog Number: MO20011 Host: Mouse

Ig Class: IgM,kappa clone: 8A11 Species Reactivity: Human

Immunogen
Sequence: Prokaryotic recombinant protein corresponding to the external

Format: Liquid- tissue culture supernatant containing 15mM sodium azide.

Sequence: corresponding to the external contains domain of the human

transforming

Applications: Immunohistochemistry-1:25-1:50 (Paraffin-embedded and frozen tissue)

Trypsin digestion of paraffin sections may enhance staining in some cases.

60 minutes primary antibody incubation at 25° C

Standard ABC technique.

Dilutions listed as a recommendation. Optimal dilution should be determined by investigator.

Storage: Antibody can be aliquotted and stored frozen at -20° C to -70° C in a manual defrost freezer for six

months without detectable loss of activity. The antibody can be stored at 2° - 8° C for 1 month without

detectable loss of activity. Avoid repeated freeze-thaw cycles.


Application Notes

Staining Pattern: Membrane (some cytoplasmic staining may be observed in certain cell types eg: lymphoid cells).

Description/Data:

Transforming growth factor beta (TGFB) inhibits cell proliferation and stimulates cell differentiation. This is achieved through a receptor complex of types I and II TGFB receptors. The type II receptor is a transmembrane serine/threonine kinase which is able to bind TGFB and transactivate the type I receptor. TGFB receptors are found in all myometrial tissues and, in the kidney, they are found exclusively in renal tubular cells. In normal gastric mucosa, type I and II receptors are demonstrated in fundic glands, but not on surface mucous cells and in normal small bowel and colon, TGFB receptors are expressed in the epithelial cells of the upper crypts.

Image: TGF Beta R (Type1) staining of paraffin embedded Human colon, ulcerative colitis. Note membrane staining of a proportion of epithelial cells and lymphocytes.



FOR RESEARCH USE ONLY

NEUROMICS' REAGENTS ARE FOR IN VITRO AND CERTAIN NON-HUMAN IN VIVO EXPERIMENTAL USE ONLY AND NOT INTENDED FOR USE IN ANY HUMAN CLINICAL INVESTIGATION, DIAGNOSIS, PROGNOSIS, OR TREATMENT. THE ABOVE ANALYSES ARE MERELY TYPICAL GUIDES. THEY ARE NOT TO BE CONSTRUED AS BEING SPECIFICATIONS. ALL OF THE ABOVE INFORMATION IS, TO THE BEST OF OUR KNOWLEDGE, TRUE AND ACCURATE. HOWEVER, SINCE THE CONDITIONS OF USE ARE BEYOND OUR CONTROL, ALL RECOMMENDATIONS OR SUGGESTIONS ARE MADE WITHOUT GUARANTEE, EXPRESS OR IMPLIED, ON OUR PART. WE DISCLAIM ALL LIABILITY IN CONNECTION WITH THE USE OF THE INFORMATION CONTAINED HEREIN OR OTHERWISE, AND ALL SUCH RSKS ARE ASSUMED BY THE USER. WE FURTHER EXPRESSLY DISCLAIM ALL WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE.-V2/08/2015